


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Images in cardio-thoracic surgery

DeBekay repair for type III thoracoabdominal aortic aneurysm

Zan Mitrev, Vladimir Belostotski, Lidiya Veljanovska, Nikola Hristov^{*}

Special Hospital for Surgery "Filip Vtori", Skopje, Macedonia

Received 30 November 2008; received in revised form 23 January 2009; accepted 29 January 2009; Available online 9 March 2009

Keywords: Aneurysm; Aortic operation

A 65-year-old man with abdominal pain, nausea and vomiting was diagnosed with thoracoabdominal aortic aneurysm, Crawford type III (Fig. 1). Immediate surgery was performed through 6th intercostal space and retro-

peritoneally, employing DeBakey repair (Fig. 2). Following repair, aneurysm sac was opened, bleeding points sutured, aneurysm neck suture ligated.



Fig. 1. Preoperative 64 slice computerized tomography of Crawford type III thoracoabdominal aortic aneurysm, maximal diameter of 10 cm.



Fig. 2. Postoperative 64 slice computerized tomography of DeBakey type repair for Crawford type III thoracoabdominal aortic aneurysm. Arrow 1 indicating proximal end to side prosthesis implantation on distal thoracic aorta; arrow 2 implantation of the celiac trunk over short 10 mm vascular graft on the prosthesis (arrow 8); arrow 3 implantation of the right renal artery over short 10 mm vascular graft on the prosthesis (arrow 11); arrow 4 implantation of the superior mesenteric artery over short 10 mm vascular graft on the prosthesis (arrow 9); arrow 5 implantation of the left renal artery over short 10 mm vascular graft on the prosthesis (arrow 10); arrow 12 end to end anastomosis between tubular and bifurcated graft; arrow 6 end to end anastomosis between prosthesis and right external iliac artery; arrow 7 end to end anastomosis between prosthesis and left external iliac artery.

^{*} Corresponding author. Address: PZU "Filip Vtori", Skopje, Macedonia.
Tel.: +389 2 3091500; fax: +389 2 3091499.
E-mail address: hristov@cardiosurgery.com.mk (N. Hristov).